sciencenewsline *nature*

Home | Top News | Most Popular | Science Video | Multimedia | News Feeds

Medicine | Nature & Earth | Biology | Technology & Engineering | Space & Planetary | Psychology | Physics & Chemistry | Economics | Archaeology

Scientists Develop New Technology to Detect Deep Sea Gas

Published: October 12, 2011. University of Southampton

A new ultra-sensitive technology which can monitor leaks from underwater gas pipelines has been developed by scientists at the University of Southampton.

The research has shown that potentially environmentally and financially disastrous gas leaks from pipelines, and methane naturally leaking from the seabed, could in future be detected using changes in acoustic signals.

- More news from University of Southampton
- University of Southampton

Using a simple set of underwater microphones to monitor these changes would provide a cost effective, unique detection system which would be one hundred times more sensitive than current monitors used by the oil and gas industry for remote detection with long deep sea pipelines.

"This new technology could save gas extraction and distribution companies millions in lost revenue. Severe leaks can also be dangerous to nearby oil rigs, shipping and for shore-based gas distribution facilities," comments Professor Tim Leighton of the University's Institute of Sound and Vibration Research who led the research.

He adds: "The technology would allow us remotely to monitor and potentially reduce the release into the atmosphere of gases from the seabed. This applies both to gas extracted by the petrochemical industries and to the methane which is naturally released from the seabed."

Natural leaks of methane gas can be damaging to the environment because it is a greenhouse pollutant.

The new acoustic technology, which is in early development, could also be used in future to monitor the structural integrity of carbon capture and storage facilities which are being developed globally. These facilities will trap carbon emissions, which scientists believe may be contributing to global warming. The UK government has just announced it is investing £1 billion in their development.

Post Comment

All comments are reviewed before being posted. We cannot accept messages that advertise a product, group or web site. If you are looking for a response to a question please use our feedback page

Related Articles »

nature Switching from Coal to Natural Gas Would Do Little for Global Climate, Study **Indicates**

Springer

Although the burning of natural gas emits far less carbon dioxide than coal, a new study concludes that a greater reliance on natural gas would fail to significantly slow down climate change. The study appears this week in the Springer journal Climatic Change Letters.




Ethane Levels Yield Information About Changes in Greenhouse Gas Emissions National Science Foundation

Recent data from NSF-funded research in both Greenland and Antarctica demonstrate that fossil-fuel related emissions of both methane and ethane, two of the most abundant hydrocarbons in the atmosphere, declined at the end of the twentieth century, according to a paper published Thursday in the journal Nature.

More Nature & Earth »

cloud



Storm Chasers of Utah: Tornadohunting Radar Truck Seeks Wasatch Snow And Rain

biodiversity



Citrus Indica Tanaka - a Progenitor Species of Cultivated Citrus

species Between Geometry And Biology: How And Why Does the Number of Species Depend on Area There are few universal rules in ecology, but arguably one is the relationship between the area of a study.

antarctic International Team to Drill Beneath Massive Antarctic Ice Shelf

> An international team of researchers funded by NASA and the National Science Foundation (NSF) will travel next month to ..

solar New Reports Urges More Detailed Utility Metering to Improve Building Efficiency Anew interagency report recommends systematic consideration of new metering technologies, called submetering. that can yield up-to-date, finely grained

methane NOAA Greenhouse Gas Index Continues to Climb NOAA's updated Annual Greenhouse Gas Index (AGGI), which measures the direct climate influence of many greenhouse gases such as .

carbon Long-Term Carbon Storage in Ganges Basin May Portend Global Warming Worsening Woods Hole Oceanographic Institution (WHOI) scientists have found that carbon is stored in the soils and sediments of the

forest Report Provides New Analysis of Carbon Accounting, Biomass Use, And Climate Benefits Arecent report provides new ideas regarding carbon and energy benefits forests and forest products provide. The report, Managing.

carbon Methane May Be Answer to 56million-year Question The release of massive amounts of carbon from methane hydrate frozen under the seafloor 56 million years

energy Americans Using More Fossil Fuels

ago has .

American energy use went back up in 2010 compared to 2009 when consumption was at a 12-year low.



nature Switching from Coal to Natural Gas Would Do Little for Global Climate, Study 9/8/11 Indicates

National Center for Atmospheric Research/University Corporation for Atmospheric Research BOULDER —Although the burning of natural gas emits far less carbon dioxide than coal, a new study concludes that a greater reliance on natural gas would fail to significantly slow down climate change.

nature Study Gives Green Light to Plants' Role in Global Warming 4/29/10 University of Edinburgh

Plants remain an effective way of tackling global warming despite emitting small amounts of an important greenhouse gas, a study has shown. Research led by the University of Edinburgh suggests that plant leaves account for less than one per cent of the Earth's emissions of methane —which is considered to be about 25 times more effective than carbon dioxide at global warming.

Greenhouse Gas Impact of Hydroelectric Reservoirs Downgraded Cary Institute of Ecosystem Studies

An international team of scientists has amassed the largest data set to date on greenhouse gas emissions from hydroelectric reservoirs. Their analysis, published today in the online version of Nature Geoscience, posits that these human-made systems emit about 1/6 of the carbon dioxide and methane previously attributed to them.

nature Methane Gas from Cows - the Proof Is in the Poo 6/7/11 University of Leeds

Scientists could have a revolutionary new way of measuring how much of the potent greenhouse gas methane is produced by cows and other ruminants, thanks to a surprising discovery in their poo. Researchers from the University of Bristol and the Teagasc Animal and Grassland Research Centre in Ireland, have found a link between methane production and levels of a compound called archaeol in the faeces of several fore-gut fermenting animals including cows, sheep and deer.

MORE»

Most Popular - Nature »

WHEAT

Using Biochar to Boost Soil Moisture

Scientists at the U.S. Department of Agriculture (USDA) are leading the way in learning more about "biochar," the charred biomass created from wood, other plant material, and manure. . . .

FOREST »



Report Provides New Analysis of Carbon Accounting, Biomass Use, And Climate Benefits

CHOLERA»



Study Predicts Sea Level Rise May Take Economic Toll on California Coast

CROPS »

ICRISAT-led Team Cracks Pigeonpea Genome

Once referred to as an "orphan crop" mainly grown by poor farmers, pigeonpea is now set to join the world's league of major food crops with the completion of ...

WEATHER »

MORE»



UA Scientists Find Evidence of Roman Period Megadrought

ScienceNewsline | Deutsch | Español | Français | Italiano | Português

Home | Top News | Most Popular | Science Video | Multimedia | News Feeds | News Search

Medicine & Health Care | Nature & Earth | Biology | Technology & Engineering | Space & Planetary | Psychology | Physics & Chemistry | Economics | Archaeology

About | Privacy Policy | Feedback